

COLON AND RECTAL SURGERY, INC.

Website: www.crscolumbus.com

K.S. Khanduja, M.D., F.A.C.S., F.A.S.C.R.S.

A. Padmanabhan, M.D., M.S., (Surg.), F.I.C.S., F.A.S.C.R.S.

C. W. Taylor III, M.D., F.A.C.S., F.A.S.C.R.S.

M.A. Lindsey, M.D., F.A.C.S.

J. Madhavan, M.D., M.S. (Surg.), F.R.C.S.Ed., F.A.C.S., F.A.S.C.R.S.

E. H. Bailey, M.D.

5965 E. Broad St., Suite 250

Columbus, OH 43213

Phone: 614-759-5060

Fax: 614-759-5065

745 W. State St., Suite 605

Columbus, OH 43222

Phone: 614-759-5060

Fax: 614-759-5065

COLONOSCOPY INSTRUCTIONS: PLEASE READ CAREFULLY

Start a clear liquid diet at breakfast **the day before** your colonoscopy. **(NO FOOD THE DAY BEFORE OR THE DAY OF)** This includes clear or strained fruit juices, jello (no red jello), clear broths, coffee (no cream or non-dairy cream), tea and other bottled beverages such as sodas and sports drinks. **No milk or milk products.**

Stop drinking 3 hours before your colonoscopy except for small sips of water to take your medications, if needed. Continue your liquid diet up until 3 hours before your colonoscopy.

You must continue your essential medications according to schedule. If you are diabetic, call your doctor who prescribes your diabetic medication to determine what dosage you need the day before your colonoscopy.

Do not take your diabetic medication the morning of your colonoscopy. You must discontinue **BLOOD THINNER** _____ days before your colonoscopy. **Please OK these changes with your Primary Care Physician or Cardiologist before stopping your BLOOD THINNER.**

Please let us know if you have had **Prosthetic Heart Valve or joint replacement surgery** within the last 6 months. You may need antibiotics before your colonoscopy.

Please look at the additional sheet for your bowel preparation instructions which must take place **the day before and the day of** your colonoscopy.

Plan to arrive at _____ AM/PM for your colonoscopy on _____ at _____ AM/PM. Your colonoscopy will be at _____ AM/PM.

Mt. Carmel East Hospital: South tower entrance behind hospital, take elevator to 2nd floor. **614-234-6000**

Mt. Carmel Medical Center: Use the main entrance valet or park in the garage close to the main entrance. From the main lobby, follow signs to the Central Elevators. Go to the 4th floor and exit left. You will be at the registration desk. Once you have registered, you will be directed to the Endoscopy Suite on the 5th floor. **614-234-5000**

Taylor Station Surgery Center: 275 Taylor Station Rd. **BUILDING B**, ½ mile north of Mt. Carmel East Hospital. **614-751-4466**

Berger Hospital (Circleville): 600 North Pickaway St. Go in the Main entrance to the registration desk. **740-474-2126**

Grove City Surgery Center: 1325 Stringtown Road, GroveCity, OH 43123. **614-991-6900**

St. Ann's Hospital: Use main entrance of the hospital. **614-898-4000.**

You will not be allowed to drive yourself home after this procedure so please arrange transportation prior to your procedure. You will be at the facility 2 to 3 hours total. You can return to your normal activities the day after your colonoscopy.

Dr. Khanduja/Dr. Padmanabhan/Dr. Taylor/Dr. Lindsey/Dr. Madhavan/Dr. Bailey will do your colonoscopy.

Colon And Rectal Surgery, Inc.

K.S. Khanduja, M.D., F.A.C.S., F.A.S.C.R.S

A. Padmanabhan, M.D., M.S., (SURG.), F.I.C.S., F.A.C.S., F.A.S.C.R.S.

C.W. Taylor III, M.D., F.A.C.S., F.A.S.C.R.S.

M.A. Lindsey, M.D., F.A.C.S.

J. Madhavan, M.D., M.S., (SURG.), F.R.C.S.Ed., F.A.C.S., F.A.S.C.R.S.

E. Bailey, M.D.

Moviprep Instructions

Moviprep kit contains: 2 pouch A packets, 2 pouch B packets & 1 disposable container

Preparation the day before your colonoscopy

START A CLEAR LIQUID DIET AT BREAKFAST the morning before your procedure. Please see the clear liquids sheet for details on clear liquids.

Mix Solution prior to 6pm:

- Combine **ONE** of the pouch A packets with **ONE** of the pouch B packets.
- Add lukewarm drinking water to the top line of the container. Mix to dissolve.
- You may refrigerate solution before drinking, must drink within 24 hours of mixing

At 6pm:

- Moviprep container is divided by 4 marked lines. Every 15 minutes, drink the solution down to the next marked line (*approximately 8 ounces*).
- Be sure to drink **ALL** of the solution (*one liter total*).
- Drink **16 ounces** of your choice of clear liquids (*see clear liquids sheet*).

At 7pm:

- Mix second dose of solution, pouch A and pouch B packets and refrigerate.

Preparation the day of your colonoscopy

- **4 hours before leaving** home for your procedure, take the second dose of solution the exact same way as the first dose.
- Drink 16 ounces of your choice of clear liquids

Tip: You may add one “on the go” crystal light raspberry lemonade packet to each dose of Moviprep to enhance the flavor

Clear Liquids

Your doctor has ordered **ONLY clear liquids the day before** your test or surgery in order to prepare your body for the procedure. **Do not eat solid foods during this time.**

*****Nothing In Red*****

You may have these liquids:

- Water
- Soft Drinks
- Gatorade and other sports drinks
- Kool-Aid (no red)
- Clear juices: apple, white cranberry, white grape, no orange juice(pulp or pulp free)
- Strained lemonade, fruit punch, powdered fruit mixes
- Tea, decaffeinated tea – no cream or non dairy creamer
- Coffee, decaffeinated coffee – no cream or non dairy creamer
- Broth, bouillon
- Jell-O (no red)
- Popsicles (no red)
- Hard candy, honey, sugar

You may NOT drink:

- Milk, smoothies or other dairy products
- Orange juice, fruit nectars or vegetable juices
- Alcohol

* If your test or surgery involves the digestive system (esophagus, stomach, or bowels) **avoid all** red liquids, red popsicles, red Jell-O and red hard candy

* **Try to limit liquids containing caffeine, as caffeine can upset an empty stomach**